HOSPITAL VETERINÁRIO
Muralha de Évora

equimuralha
Medicina Veterinária Equina

COM O APOIO DE:

HIPRA

14

**JORNADAS
INTERNACIONAIS**
HOSPITAL VETERINÁRIO
MURALHA DE ÉVORA

PRODUTOS LÁCTEOS OBTIDOS
COM LEITE DE PEQUENOS
RUMINANTES: DESAFIOS E
OPORTUNIDADES PARA O
SETOR

Carlos Dias Pereira

Escola Superior
Agrária de Coimbra

SUMÁRIO

1-A problemática dos queijos de ovelha e cabra portugueses com DOP;

2-Valorização de soro e de soroelho: oportunidades para o desenvolvimento de novos produtos;

3-Manteiga e leite de ovelha com adição de microrganismos probióticos;

4-Recuperação de energia no processo de fabrico de requeijão.

A problemática dos queijos de ovelha e cabra portugueses com DOP

Os queijos com DOP representam uma parcela mínima dos queijos de ovelha e cabra produzidos em Portugal (*ca.* 9% e 1% respetivamente).

QUAIS AS CAUSAS DESTA SITUAÇÃO?

Atualmente não falta informação de excelência sobre as características físico-químicas e microbiológicas dos queijos DOP Portugueses (especialmente os produzidos com queijo de ovelha).

QUAIS AS RAZÕES PARA O CONHECIMENTO CIENTÍFICO NÃO SER TRANSFERIDO PARA OS PRODUTORES?

Article

Serra da Estrela PDO Cheese Microbiome as Revealed by Next Generation Sequencing

Rui Rocha * , Manuela Vaz Velho , Joana Santos and Paulo Fernandes

Microorganisms **2021**, *9*, 2007. <https://doi.org/10.3390/microorganisms9102007>

Figure 9. Venn diagram representing shared core taxa of bacteria between raw ewes' milk (blue area), cardoon (red area) and Serra da Estrela cheese (green area) matrices. f denotes identified bacterial families without taxonomic attribution to genus and/or species.

Figure 10. Venn diagram representing shared core taxa of fungi between raw ewes' milk (blue area), cardoon (red area) and Serra da Estrela cheese (green area) matrices. f denotes identified bacterial families without taxonomic attribution to genus and/or species.

Bacterial Communities in Serpa Cheese by Culture Dependent Techniques, 16S rRNA Gene Sequencing and High-throughput Sequencing Analysis

Maria Teresa P. Gonçalves, María José Benito , María de Guía Córdoba, Conceição Egas, Almudena V. Merchán , Ana I. Galván, and Santiago Ruiz-Moyano

Vol. 83, Iss. 5, 2018 • Journal of Food Science

<https://www.serpaflores.com/>

Figure 1–Mean relative abundance of bacteria in cheeses from different industries (capital letters) analyzed in two different seasons (winter and spring) using high-throughput sequencing of the 16S rRNA gene. (A) at the family level; (B) at the genus level.

Article

Impact of LAB from Serpa PDO Cheese in Cheese Models: Towards the Development of an Autochthonous Starter Culture

Helena Araújo-Rodrigues ¹ , António P. L. Martins ^{2,3,†} , Freni K. Tavoria ¹ , João Dias ^{3,4} ,
Maria Teresa Santos ⁴ , Nuno Alvarenga ^{2,3} and Manuela E. Pintado ^{1,*}

Valorização de soro e de soroelho: oportunidades para o desenvolvimento de novos produtos

O soro e o soroelho representam um dos principais desafios/opportunidades do setor dos laticínios.

Em Portugal são produzidas anualmente cerca de 600.000 toneladas de soro, na sua maioria resultantes da produção de queijos de leite de vaca (65.736 ton queijo).

No entanto, a produção de soro e de soroelho resultante do fabrico de queijos de pequenos ruminantes não é negligenciável (11.870 ton queijos de ovelha; 4.221 queijos de cabra e 6.160 ton queijos de mistura).

No caso do soro de queijos de ovelha e cabra, uma parte é utilizada na produção de requeijão.

No entanto, devido aos custos energéticos do processo de fabrico de requeijão e ao curto prazo de validade deste produto, uma parte significativa do soro não é processada.

Por outro lado, do fabrico de requeijão resulta o soroelho que não é aproveitado em alimentação humana e representa um problema ambiental, se descartado sem tratamento adequado.

Valorização de soro e
de soroelho:
oportunidades para o
desenvolvimento de
novos produtos

PROCESSOS DE FILTRAÇÃO TANGENCIAL

- Processos de filtração seletiva por membrana sob ação de um gradiente de pressão.
- O líquido de alimentação (A), que circula paralelamente à membrana, é impulsionado por uma bomba que cria o gradiente de pressão (Δp) suficiente para que o filtrado ou permeado (Pe) a atravesse.
- O concentrado ou retido (Re) corresponde à porção de (A) que não atravessa a membrana e, na qual, um ou vários dos seus componentes aumentam de concentração.
- O fluxo de filtração corresponde ao caudal de permeado que passa através de determinada área ($Lh^{-1} m^{-2}$).

Figure 7.10. Batch ultrafiltration with partial recycle of retentate. If feed is added continuously at the same rate as the flux, this is the "batch with topped off" mode.

CONVENTIONAL FILTRATION

TANGENTIAL FILTRATION

MICROFILTRATION

SUSPENDED PARTICLES,
BACTERIA, CELLS

ULTRAFILTRATION

MACROMOLECULES

NANOFILTRATION

SUGARS
DIVALENT SALTS
DISSOCIATED ACIDS

REVERSE OSMOSIS

MONOVALENT SALTS
UNDISSOCIATED ACIDS

WATER

Fermented dairy products based on ovine cheese whey

C. D. Pereira¹ · M. Henriques¹ · D. Gomes¹ · R. Gouveia¹ · A. Gomez-Zavaglia² ·
G. de Antoni^{2,3}

Fig. 1 Gross chemical composition of: **a** liquid-type products fermented by kefir grains (K1), commercial mix of probiotic (P1) and kefir grains+commercial mix of probiotic (K+P1); **b** gel-type products fermented by kefir grains (K2), commercial mix of probiotic (P2) and kefir grains+commercial mix of probiotic (K+P2). Total Solids (TS). Different letters (**a**, **b**...) indicate statistically significant differences ($P < 0.05$) within fermented products for each formulation

Fig. 8 Sensorial evaluation of liquid-type fermented products (K1, P1 and K+P1) and gel-type fermented products (K2, P2 and K+P2)

Article

Sheep's and Goat's Frozen Yoghurts Produced with Ultrafiltrated Whey Concentrates

Natalí Garcia Marnotes ^{1,2}, Arona Figueroa Pires ¹, Olga Díaz ², Angel Cobos ² and Carlos Dias Pereira ^{1,3,*}

Appl. Sci. **2021**, *11*, 6568. <https://doi.org/10.3390/app11146568>

Table 1. Formulations used for the production of frozen yoghurts (% *w/w*).

Product	0% Inulin	2.5% Inulin	5% Inulin
LWC yoghurt	89.4	86.9	84.4
Inulin ¹	0.0	2.5	5.0
Guar gum ²	0.1	0.1	0.1
Xanthan gum ³	0.1	0.1	0.1
Honey ⁴	5.0	5.0	5.0
Sugar ⁵	5.0	5.0	5.0
Citric acid ⁶	0.4	0.4	0.4

Suppliers of ingredients: ¹ Inulin: Fibruline™XL, Cosucra, supplied by Induxtra de Suministros, Moita, Portugal; ^{2,3} Formulab, Maia, Portugal; ⁴ Aldi, Portugal; ⁵ RAR, Porto, Portugal; ⁶ J.M. Vaz Pereira, Lisbon, Portugal.

Figure 1. Gross composition of ewe's and goat's yoghurts produced with UF concentrated cheese whey. Different letters indicate significant differences ($p < 0.05$) between components of ewe's and goat's yoghurts.

Table 2. Texture parameters of ewe's and goat's yoghurts one day after production (mean values and standard deviation).

Parameter	Ewe's Yoghurt		Goat's Yoghurt	
	Mean	±	Mean	±
Hardness (N)	13.55 ^a	0.33	0.50 ^b	0.08
Adhesiveness (N.s)	0.53 ^a	0.03	-0.51 ^b	0.10
Gumminess (N)	13.52 ^a	0.34	0.47 ^b	0.08
Cohesiveness	1.00 ^a	0.01	0.93 ^b	0.02
Resilience	0.90 ^a	0.02	0.04 ^b	0.01
Viscosity (Cps)	16 946.67 ^a	46.19	143.33 ^b	12.66

Different superscript letters indicate significant ($p < 0.05$) differences between values in rows.

Figure 2. Elastic (G') and viscous (G'') moduli and complex viscosity of ewe's (A,B) and goat's (C,D) yoghurts produced with UF concentrated cheese whey.

Figure 4. Gross chemical composition of ewe's and goat's frozen yoghurts produced with UF concentrated cheese whey over 21 days of refrigerated storage (EW. 0% = ewe's yoghurt with 0% inulin addition; EW. 2.5% = ewe's yoghurt with 2.5% inulin addition; EW. 5.0% = ewe's yoghurt with 5% inulin addition; GO. 0% = goat's yoghurt with 0% inulin addition; GO. 2.5% = goat's yoghurt with 2.5% inulin addition; GO. 5% = goat's yoghurt with 5% inulin addition). Different letters indicate significant differences ($p < 0.05$) between components of ewe's and goat's frozen yoghurt formulations.

Table 5. Overrun and meltdown rate of ewe's and goat's frozen yoghurts immediately after production ($n = 3$).

	EW. 0%	$\pm$	EW. 2.5%	$\pm$	EW. 5.0%	$\pm$	GO. 0%	$\pm$	GO. 2.5%	$\pm$	GO. 5.0%	$\pm$
Overrun	31.04 ^a	1.89	31.09 ^a	2.04	40.29 ^b	2.20	29.34 ^a	1.45	31.13 ^a	1.56	29.34 ^a	2.23
Meltdown rate	3.47 ^a	0.03	6.33 ^b	0.03	4.41 ^c	0.10	27.99 ^d	0.01	13.07 ^e	0.50	15.00 ^f	0.55

EW. 0% = ewe's yoghurt with 0% inulin addition; EW. 2.5% = ewe's yoghurt with 2.5% inulin addition; EW. 5.0% = ewe's yoghurt with 0% inulin addition; GO. 0% = goat's yoghurt with 0% inulin addition; GO. 2.5% = goat's yoghurt with 2.5% inulin addition; GO. 5.0% = goat's yoghurt with 5% inulin addition. Different superscript letters (a,b, ...) indicate significant ($p < 0.05$) differences between values in rows.

Figure 7. *Lactobacilli* sp. (*Lb.*) and *Streptococci* sp. (*St.*) counts (log CFU g⁻¹) of ewe's and goat's frozen yoghurts produced with UF concentrated cheese whey at the day of production (EW. 0% = ewe's yoghurt with 0% inulin addition; EW. 2.5% = ewe's yoghurt with 2.5% inulin addition; EW. 5.0% = ewe's yoghurt with 5% inulin addition; GO. 0% = goat's yoghurt with 0% inulin addition; GO. 2.5% = goat's yoghurt with 2.5% inulin addition; GO. 5% = goat's yoghurt with 5% inulin addition). Different superscript letters indicate significant differences ($p < 0.05$) between products for lactobacilli (a, b, c) and streptococci (A, B, C) counts.

Table 7. Microbial counts (log CFU g⁻¹) of ewe's and goat's frozen yoghurt over storage. Lb. = lactobacilli; St. = streptococci (mean values and standard deviation).

Days		0% Inulin		2.5% Inulin		5.0% Inulin	
Ewe's		Mean	±	Mean	±	Mean	±
1	Lb.	6.50 ^a	0.71	7.02 ^a	1.09	6.76 ^a	0.59
	St.	9.11 ^a	0.45	9.05 ^a	0.33	8.82 ^a	0.28
7	Lb.	7.86 ^a	0.41	7.92 ^a	0.39	7.88 ^a	0.32
	St.	9.28 ^a	0.94	9.07 ^a	0.54	8.56 ^a	0.28
14	Lb.	6.00 ^a	1.41	6.05 ^a	0.02	5.80 ^a	0.30
	St.	8.22 ^a	0.37	8.01 ^a	0.49	8.09 ^a	0.62
21	Lb.	5.33 ^a	0.47	5.86 ^a	1.26	5.00 ^a	0.10
	St.	0.00 ^a	0.00	0.00 ^a	0.00	0.00 ^a	0.00
Goat's		Mean	±	Mean	±	Mean	±
1	Lb.	7.02 ^a	0.66	8.14 ^{ab}	1.84	5.50 ^b	0.70
	St.	7.48 ^a	1.53	5.42 ^a	0.60	5.44 ^b	0.61
7	Lb.	6.99 ^a	0.97	0.00 ^b	0.00	6.14 ^a	0.63
	St.	5.65 ^a	0.49	6.68 ^a	1.23	7.85 ^a	1.58
14	Lb.	0.00 ^a	0.00	0.00 ^a	0.00	0.00 ^a	0.00
	St.	6.94 ^a	0.11	5.43 ^b	0.51	5.74 ^b	0.37
21	Lb.	0.00 ^a	0.00	0.00 ^a	0.00	0.00 ^a	0.00
	St.	6.15 ^a	1.63	5.33 ^a	0.58	6.92 ^a	0.99

Different superscript letters (a,b) indicate significant ($p < 0.05$) differences between values in rows.

Figure 8. Sensory scores of ewe's and goat's yoghurts and correspondent frozen yoghurts produced with UF concentrated cheese whey ((A) = ewe's and goat's yoghurts; (B) = ewe's frozen yoghurts, (C) = goat's frozen yoghurts).

Article

Evaluation of the Characteristics of Sheep's and Goat's Ice Cream, Produced with UF Concentrated Second Cheese Whey and Different Starter Cultures

Arona Pires ^{1,2}, David Gomes ¹, João Noronha ¹ , Olga Díaz ², Angel Cobos ² and Carlos Dias Pereira ^{1,3,*}

Figure 1. Chemical composition of sheep's and goat's second cheese whey ice creams produced with different starter cultures. Different letters of the same component indicate significant differences ($p < 0.05$) among ice creams.

Figure 4. Complex viscosity (η^*) of sheep's and goat's second cheese whey ice creams produced with different starter cultures over storage. 1 day (A); 30 days (B); 60 days (C); 120 days (D).

Figure 6. Evaluation of microbial counts of sheep's and goat's second cheese whey ice creams produced with different starter cultures over storage. *Lactobacilli* sp. (A); *Lactococci* sp. (B). Different lowercase letters in each type of ice cream indicate significant differences ($p < 0.05$) among storage days and different capital letters in each storage day indicate significant differences ($p < 0.05$) among ice creams.

Figure 7. Sensory scores of sheep's and goat's second cheese whey ice creams produced with different starter cultures over 120 days of storage. Color (A); Aroma (B); Taste (C); Texture (D).

Article

Development and Characterization of a Novel Sustainable Probiotic Goat Whey Cheese Containing Second Cheese Whey Powder and Stabilized with Thyme Essential Oil and Sodium Citrate

Manuel Mântua Esteves Garcia ¹, Carlos José Dias Pereira ² , Ana Cristina Freitas ¹ ,
Ana Maria Pereira Gomes ¹ and Maria Manuela Estevez Pintado ^{1,*}

Foods **2022**, *11*, 2698. <https://doi.org/10.3390/foods11172698>

Table 4. Physicochemical composition of whey cheeses matrices (average $\pm$ standard deviation).

Parameter	Whey Cheese Matrix				
	WC	WCP	WCPS	WCPST	WCPSTC
Dry matter (%)	27.02 $\pm$ 0.11 ^a	25.17 $\pm$ 0.04 ^b	27.35 $\pm$ 0.04 ^a	25.92 $\pm$ 0.20 ^c	25.64 $\pm$ 0.13 ^{bc}
Ash (%)	3.32 $\pm$ 0.07 ^a	3.33 $\pm$ 0.01 ^a	3.08 $\pm$ 0.08 ^{ab}	2.48 $\pm$ 0.01 ^c	2.81 $\pm$ 0.02 ^b
Fat (%)	6.13 $\pm$ 0.31 ^a	5.71 $\pm$ 0.10 ^b	6.45 $\pm$ 0.31 ^a	8.03 $\pm$ 0.09 ^c	7.59 $\pm$ 0.03 ^c
Protein (%)	11.10 $\pm$ 1.06 ^{ab}	10.84 $\pm$ 0.14 ^a	12.66 $\pm$ 0.08 ^b	11.26 $\pm$ 0.25 ^{ab}	10.78 $\pm$ 0.08 ^a

WC (non-inoculated); WCP (control); WCPS (with SCWP); WCPST (with SCWP and thyme EO); WCPSTC (with SCWP, thyme EO and sodium citrate). ^{a,b,c} Means within the same line labeled with different superscripts differ significantly ($p < 0.05$).

Table 3. Evolution of viable counts (log CFU g⁻¹) and physicochemical parameters throughout storage period (average ± standard deviation).

Strain	Storage Time (Day)	Whey Cheese Matrix			
		WCP	WCPS	WCPST	WCPSTC
<i>Bifidobacterium animalis</i> subsp. <i>lactis</i> BB-12	0	8.19 ± 0.10 ^a	8.30 ± 0.18 ^a	7.24 ± 0.07 ^b	7.38 ± 0.27 ^b
	7	8.40 ± 0.17 ^a	8.39 ± 0.13 ^a	7.24 ± 0.07 ^b	7.11 ± 0.26 ^b
	14	8.78 ± 0.08 ^a	8.99 ± 0.44 ^a	7.36 ± 0.05 ^b	7.31 ± 0.05 ^b
	21	8.93 ± 0.07 ^a	8.69 ± 0.19 ^a	7.64 ± 0.11 ^b	7.45 ± 0.05 ^b
<i>Lactobacillus rhamnosus</i> Rosell-11	0	7.99 ± 0.05 ^a	8.24 ± 0.19 ^a	7.25 ± 0.09 ^b	7.20 ± 0.10 ^b
	7	8.46 ± 0.26 ^a	8.36 ± 0.09 ^a	7.31 ± 0.09 ^b	7.26 ± 0.10 ^b
	14	8.64 ± 0.18 ^a	8.28 ± 0.17 ^a	7.34 ± 0.07 ^b	7.35 ± 0.13 ^b
	21	8.71 ± 0.15 ^a	8.46 ± 0.14 ^{ab}	7.98 ± 0.11 ^c	8.06 ± 0.10 ^{bc}
pH	0	5.72 ± 0.01 ^a	5.66 ± 0.02 ^a	6.02 ± 0.01 ^b	6.25 ± 0.00 ^c
	7	5.59 ± 0.01 ^a	5.49 ± 0.04 ^a	5.87 ± 0.01 ^b	6.21 ± 0.02 ^c
	14	4.80 ± 0.02 ^a	5.06 ± 0.08 ^b	5.39 ± 0.01 ^c	5.57 ± 0.07 ^d
	21	4.79 ± 0.00 ^a	4.87 ± 0.08 ^a	4.77 ± 0.01 ^a	5.05 ± 0.01 ^b
Titratable acidity (% Lactic acid)	0	0.42 ± 0.00 ^a	0.52 ± 0.03 ^a	0.49 ± 0.01 ^a	0.43 ± 0.01 ^a
	7	0.55 ± 0.02 ^a	0.65 ± 0.04 ^a	0.64 ± 0.02 ^a	0.53 ± 0.03 ^a
	14	0.93 ± 0.02 ^a	0.84 ± 0.07 ^{ab}	0.83 ± 0.03 ^{ab}	0.79 ± 0.03 ^b
	21	1.10 ± 0.07 ^{ab}	1.17 ± 0.10 ^{ab}	1.24 ± 0.01 ^b	1.07 ± 0.01 ^a

WCP (control); WCPS (with SCWP); WCPST (with SCWP and thyme EO); WCPSTC (with SCWP, thyme EO and sodium citrate). ^{a,b,c,d} Means within the same line labeled with different superscripts differ significantly ($p < 0.05$).

Manteiga e leitelho de ovelha com adição de microrganismos probióticos

A produção de produtos lácteos fermentados com base em leite de pequenos ruminantes (*e.g.* iogurtes) pode também ser uma oportunidade de diversificação do portfolio de produtos das empresas.

Podem ainda ser obtidos outros produtos, como sejam a manteiga de ovelha/cabra e o leitelho fermentado.

Article

Sheep's Butter and Correspondent Buttermilk Produced with Sweet Cream and Cream Fermented by Aromatic Starter, Kefir and Probiotic Culture

Tânia Silva ¹, Arona Pires ¹, David Gomes ¹, Jorge Viegas ¹, Susana Pereira-Dias ^{1,2}, Manuela E. Pintado ³,
Marta Henriques ^{1,2} and Carlos Dias Pereira ^{1,2,*}

Foods **2023**, *12*, 331. <https://doi.org/10.3390/foods12020331>

Table 1. Chemical composition of sheep's butters produced with sweet cream and with cream fermented with different starter cultures. Different superscript lowercase letters in each column (a,b) indicate significant differences ($p < 0.05$) between butters.

Products	Dry Matter (% <i>w/w</i>)	Fat (% <i>w/w</i>)	Protein (% <i>w/w</i>)	Ashes (% <i>w/w</i>)
S—Sweet	87.24 ± 2.79 ^a	83.63 ± 3.28 ^a	0.34 ± 0.05 ^a	1.19 ± 0.23 ^a
A—Aromatic	87.57 ± 5.02 ^a	83.42 ± 4.96 ^a	1.08 ± 0.14 ^b	1.35 ± 0.27 ^a
K—Kefir	85.58 ± 4.92 ^a	81.88 ± 3.89 ^a	0.84 ± 0.19 ^b	1.38 ± 0.23 ^a
P—Probiotics	78.50 ± 3.13 ^b	74.75 ± 2.18 ^b	0.89 ± 0.11 ^b	1.58 ± 0.26 ^a

Figure 1. Evolution of pH (A) and titratable acidity (B) of sheep's butters produced with sweet cream and with cream fermented with different starter cultures. Different lowercase letters (a,b,c,d) indicate significant differences ($p < 0.05$) between days of storage; Different capital letters (A,B,C,D) indicate significant differences ($p < 0.05$) between products at the same storage day.

Table 4. Texture parameters of sheep's butters produced with sweet cream and with cream fermented with different starter cultures. Average values and standard deviation ($\pm$). Different superscript lowercase letters in each column (a,b) indicate significant differences ($p < 0.05$) between products.

Products	Hardness (N)	Adhesiveness (N)	Gumminess (N)	Springiness	Cohesiveness	Resilience
S—Sweet	38.75 $\pm$ 7.30 ^a	-7.16 $\pm$ 2.39 ^a	10.80 $\pm$ 3.24 ^a	0.85 $\pm$ 0.10 ^a	0.29 $\pm$ 0.02 ^a	0.09 $\pm$ 0.02 ^a
A—Aromatic	30.78 $\pm$ 2.89 ^b	-7.98 $\pm$ 1.81 ^a	10.15 $\pm$ 1.77 ^a	0.78 $\pm$ 0.12 ^a	0.34 $\pm$ 0.04 ^a	0.09 $\pm$ 0.01 ^a
K—Kefir	37.52 $\pm$ 5.17 ^a	-6.97 $\pm$ 1.00 ^a	11.24 $\pm$ 2.15 ^a	0.80 $\pm$ 0.08 ^a	0.29 $\pm$ 0.02 ^a	0.08 $\pm$ 0.02 ^a
P—Probiotics	43.78 $\pm$ 12.43 ^a	-5.46 $\pm$ 2.32 ^a	12.66 $\pm$ 3.00 ^a	0.73 $\pm$ 0.19 ^a	0.27 $\pm$ 0.03 ^a	0.10 $\pm$ 0.01 ^a

Table 5. Fatty acid profiles of sheep's butters produced with sweet cream and with cream fermented with different starter cultures. Average values and standard deviation. Different superscript lowercase letters in each row (a,b,c) indicate significant differences ($p < 0.05$) between products.

Fatty Acids	S—Sweet	A—Aromatic	K—Kefir	P—Probiotics
C6:0	0.66 ± 0.02 ^a	0.66 ± 0.00 ^a	0.55 ± 0.00 ^b	0.79 ± 0.00 ^c
C8:0	0.88 ± 0.03 ^a	0.87 ± 0.00 ^a	0.77 ± 0.00 ^b	1.07 ± 0.00 ^c
C10:0	3.85 ± 0.11 ^a	3.78 ± 0.01 ^a	3.53 ± 0.00 ^b	4.54 ± 0.00 ^c
C12:0	3.28 ± 0.09 ^a	3.22 ± 0.00 ^a	3.13 ± 0.01 ^b	3.69 ± 0.01 ^c
C14:0	10.67 ± 0.29 ^a	10.52 ± 0.01 ^a	10.44 ± 0.05 ^a	11.39 ± 0.03 ^b
C14:1	0.19 ± 0.01 ^a	0.18 ± 0.00 ^a	0.18 ± 0.00 ^a	0.19 ± 0.00 ^a
C15:0	1.23 ± 0.02 ^a	1.20 ± 0.02 ^a	1.21 ± 0.01 ^a	1.27 ± 0.00 ^a
C15:1 c10	0.34 ± 0.00 ^a	0.33 ± 0.00 ^a	0.33 ± 0.00 ^a	0.34 ± 0.02 ^a
C16:0	32.24 ± 0.10 ^a	31.60 ± 0.10 ^a	31.80 ± 0.19 ^a	32.53 ± 0.05 ^a
C16:1 c7	0.26 ± 0.00 ^a	0.26 ± 0.00 ^a	0.26 ± 0.00 ^a	0.26 ± 0.01 ^a
C16:1 c9	1.22 ± 0.01 ^a	1.20 ± 0.01 ^a	1.21 ± 0.01 ^a	1.27 ± 0.00 ^a
C17:0	0.84 ± 0.03 ^a	0.82 ± 0.00 ^a	0.42 ± 0.57 ^b	0.83 ± 0.00 ^a
C17:1 c10	0.31 ± 0.01 ^a	0.30 ± 0.00 ^a	0.30 ± 0.00 ^a	0.33 ± 0.02 ^a
C18:0	10.56 ± 0.34 ^a	10.25 ± 0.04 ^a	10.39 ± 0.07 ^a	10.18 ± 0.01 ^a
C18:1 c9	25.47 ± 2.17 ^{ab}	26.73 ± 0.09 ^a	27.50 ± 0.18 ^a	23.16 ± 0.12 ^b
C18:1 c11	0.69 ± 0.01 ^a	0.67 ± 0.00 ^a	0.69 ± 0.00 ^a	0.66 ± 0.00 ^a
C18:1 c12	0.42 ± 0.02 ^a	0.43 ± 0.00 ^a	0.44 ± 0.00 ^a	0.40 ± 0.00 ^b
C18:2c9t12	0.42 ± 0.01 ^a	0.41 ± 0.00 ^a	0.41 ± 0.00 ^a	0.44 ± 0.00 ^b
C18:2	5.17 ± 0.02 ^a	5.12 ± 0.02 ^a	5.17 ± 0.04 ^a	5.36 ± 0.02 ^b
C18:3 c6c9c13	0.48 ± 0.01 ^a	0.41 ± 0.12 ^a	0.48 ± 0.01 ^a	0.53 ± 0.00 ^a
C18:3	0.06 ± 0.00 ^a	0.06 ± 0.00 ^a	0.07 ± 0.00 ^a	0.06 ± 0.00 ^a
C20:1	0.34 ± 0.01 ^a	0.60 ± 0.39 ^a	0.33 ± 0.00 ^a	0.32 ± 0.00 ^a
C20:3	0.15 ± 0.00 ^a	0.14 ± 0.00 ^b	0.14 ± 0.00 ^b	0.13 ± 0.00 ^c
C21:0	0.08 ± 0.00 ^a	0.08 ± 0.00 ^a	0.08 ± 0.00 ^a	0.07 ± 0.00 ^b
C22:0	0.05 ± 0.00 ^a	0.05 ± 0.00 ^a	0.05 ± 0.00 ^a	0.05 ± 0.00 ^a
C22:2	0.07 ± 0.00 ^a	0.06 ± 0.00 ^b	0.06 ± 0.00 ^b	0.06 ± 0.00 ^b
C24:0	0.05 ± 0.02 ^a	0.05 ± 0.00 ^a	0.05 ± 0.00 ^a	0.05 ± 0.02 ^a
C24:1	0.03 ± 0.00 ^a	0.01 ± 0.02 ^a	0.04 ± 0.00 ^a	0.02 ± 0.00 ^a
ΣSFA	64.38 ± 1.92 ^a	63.09 ± 0.14 ^a	62.38 ± 0.024 ^a	66.46 ± 0.12 ^b
ΣMUFA	29.26 ± 0.27 ^a	30.71 ± 0.27 ^a	31.28 ± 0.20 ^a	26.96 ± 0.11 ^b
ΣPUFA	6.36 ± 0.20 ^a	6.19 ± 0.14 ^a	6.34 ± 0.05 ^a	6.58 ± 0.01 ^b

SFA = saturated fatty acids; MUFA = monounsaturated fatty acids; PUFA = polyunsaturated fatty acids.

Figure 2. Rheological parameters of sheep's butters produced with sweet cream (S) and cream fermented with aromatic starter culture (A), kefir (K) and probiotics (P). **(A)** Elastic (G') and viscous (G'') moduli; **(B)** complex viscosity. According to power law model (Equation (2)) consistency index (K) and power law index (n).

Table 6. Physicochemical characteristics of sheep's buttermilks produced with sweet cream and with cream fermented with different starter cultures. Different superscript lowercase letters in each row (a,b,c,d) indicate significant differences ($p < 0.05$) between products. (T.A. = titratable acidity).

Parameters	Products			
	S—Sweet	A—Aromatic	K—Kefir	P—Probiotics
Total solids (% <i>w/w</i>)	13.92 ± 0.13 ^a	11.87 ± 0.11 ^b	13.70 ± 0.11 ^c	10.49 ± 0.02 ^d
Protein (% <i>w/w</i>)	5.12 ± 0.00 ^a	4.25 ± 0.10 ^b	3.70 ± 0.07 ^c	3.95 ± 0.05 ^d
Fat (% <i>w/v</i>)	2.57 ± 0.06 ^a	2.53 ± 0.12 ^a	2.93 ± 0.58 ^a	2.10 ± 0.00 ^b
Ashes (% <i>w/w</i>)	1.52 ± 0.11 ^a	0.68 ± 0.07 ^b	0.78 ± 0.33 ^b	0.76 ± 0.10 ^b
T.A. (mL NaOH 1 N/L)	35.10 ± 0.47 ^a	84.98 ± 0.64 ^b	56.3 ± 0.49 ^c	84.75 ± 0.90 ^b
pH	6.24 ± 0.01 ^a	4.65 ± 0.00 ^b	4.92 ± 0.01 ^c	4.52 ± 0.01 ^d
Viscosity (cPs)	18.42 ± 1.42 ^a	43.08 ± 4.31 ^b	165.77 ± 1.97 ^c	219.87 ± 8.81 ^d
Color values				
L*	82.07 ± 0.15 ^a	77.00 ± 0.35 ^b	83.70 ± 0.10 ^c	82.40 ± 0.26 ^a
a*	-3.97 ± 0.15 ^a	-3.57 ± 0.25 ^b	-3.83 ± 0.06 ^b	-4.20 ± 0.10 ^c
b*	4.27 ± 0.21 ^a	0.97 ± 0.06 ^b	5.97 ± 0.06 ^c	5.33 ± 0.21 ^d

Figure 3. Microbiological characteristics of sheep's butters and correspondent buttermilks produced with sweet cream and with cream fermented with different starter cultures. (A) lactobacilli in butter; (B) lactococci in butter; (C) yeasts and molds in butter; (D) lactobacilli in buttermilk; (E) lactococci in buttermilk; (F) yeasts and molds in buttermilk; different lowercase letters (a,b,c,d) indicate significant differences ($p < 0.05$) between days of storage; different capital letters (A,B,C,D) indicate significant differences ($p < 0.05$) between products at the same storage day.

Pedidos - BB/CA1K

A publicação dos pedidos de modelos de utilidade a seguir indicados é efetuada nos termos do disposto no artigo 131.º do Código da Propriedade Industrial; da data de publicação do presente aviso começa a contar-se o prazo de dois meses para a apresentação de reclamações de quem se julgar prejudicado pela eventual concessão dos mesmos, nos termos do artigo 17.º do mesmo Código.

Recuperação de energia no processo de fabrico de requeijão

- (11) **12150** (13) U
 (22) 2020.06.30
 (30)
 (71) PT INSTITUTO POLITÉCNICO DE
 COIMBRA
 (72) CARLOS JOSÉ DIAS PEREIRA
 DAVID MANUEL GAMA SIMÕES GOMES
 ARONA FIGUEROA PIRES
 MARTA HELENA FERNANDES HENRIQUES
 NATALÍ GARCÍA MARNOTES
 JORGE MANUEL PRATAS DOS SANTOS
 AREDE
 JORGE FERREIRA VIEGAS
 (51) **Int. Cl.**
A01J 25/00 (2006.01) A01J 25/11 (2006.01)
F28D 7/00 (2006.01) F04B 43/12 (2006.01)
 (54) **SISTEMA AUTOMATIZADO DE**
PRODUÇÃO DE REQUEIJÃO COM
RECUPERAÇÃO DE CALOR

(28)
 (57) SISTEMA PARA PRODUÇÃO DE REQUEIJÃO COM RECUPERARÃO DE CALOR CARACTERIZADO POR COMPREENDER UM TANQUE DE BALANÇO (1); SENSORES DE NÍVEL(2, 7, 16, 17); UMA BOMBA PERISTÁLTICA (3); UM PERMUTADOR DE CALOR (4) PARA TRANSFERIR CALOR ENTRE FLUIDOS QUE CIRCULAM EM CONTRACORRENTE; PELO MENOS UM TANQUE DE PRÉ-AQUECIMENTO (5, 8) DO SORO RESPETIVAMENTE COM UMA SERPENTINA (6, 9) TUBULAR PARA A CIRCULAÇÃO DE SORELHO; TRÊS ELECTROVÁLVULAS (10, 11, 12); UM TANQUE DE FABRICO DE REQUEIJÃO (13) COM UM AGITADOR DE PÁ (14) E MOTOR ELÉTRICO ADAPTÁVEL AO TOPO DO TANQUE DE FABRICO DE REQUEIJÃO (13); UMA VÁLVULA DE ADMISSÃO DE VAPOR (15); UMA VÁLVULA DE SEGURANÇA; UMA BOMBA CENTRIFUGA (19) PARA CIRCULAÇÃO DO SORELHO ATRAVÉS DA PELO MENOS UMA SERPENTINA (6, 9) E DO PERMUTADOR DE CALOR (4); UM PAINEL DE CONTROLO (20); UMA PLATAFORMA DE SUPORTE (21); UM TANQUE AUXILIAR DE SORELHO (TA1) E UM TANQUE AUXILIAR DE SORO (TA2).

Figura 1

[Ver Fascículo Completo](#)

- 1- TANQUE DE BALANÇO
- 2- SENSOR DE NÍVEL 1
- 3- BOMBA PERISTÁLTICA PARA SORO
- 4- PERMUTADOR DE CALOR
- 5- SENSOR DE NÍVEL 2
- 6- TANQUE DE PRÉAQUECIMENTO DE SORO 1
- 7- SERPENTINA AMOVÍVEL 1
- 8- ELECTROVÁLVULA 1
- 9- TANQUE DE PRÉAQUECIMENTO DE SORO 2
- 10- SERPENTINA AMOVÍVEL 2

- 11- ELECTROVÁLVULA 2
- 12- ELECTROVÁLVULA 3
- 13- AGITADOR DE PÁ
- 14- SENSOR DE NÍVEL 3
- 15- VÁLVULA DE ADMISSÃO DE VAPOR
- 16- SENSOR DE NÍVEL 4
- 17- TANQUE DE FABRICO DE REQUEIJÃO
- 18- VÁLVULA DE SEGURANÇA
- 19- BOMBA DE CIRCULAÇÃO DE SORELHO
- 20- QUADRO DE CONTROLO

- 1- TANQUE DE BALANÇO
- 2- SENSOR DE NÍVEL 1
- 3- BOMBA PERISTÁLTICA PARA SORO
- 4- PERMUTADOR DE CALOR (1 E 2)
- 5- TANQUE DE PRÉAQUECIMENTO DE SORO
- 6- SERPENTINA AMOVÍVEL
- 7- SENSOR DE NÍVEL 2
- 10- ELETROVÁLVULA
- 13- TANQUE DE FABRICO DE REQUEIJÃO
- 14- AGITADOR DE PÁ

- 15- VÁLVULA DE ADMISSÃO DE VAPOR
- 16- SENSOR DE NÍVEL 3
- 17- SENSOR DE NÍVEL 4
- 18- VÁLVULA DE SEGURANÇA
- 19- BOMBA DE CIRCULAÇÃO DE SORELHO
- 20- QUADRO DE CONTROLO
- 21- PLATAFORMA DE APOIO
- 22- BOMBA DE CIRCULAÇÃO DE ÁGUA
- 23- TERMOACUMULADOR

Escola Superior Agrária

Politécnico de Coimbra

OBRIGADO PELA
VOSSA ATENÇÃO

www.lacties.com

LACTIES

Inovação, Eco-Eficiência e Segurança